



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by ANNA L. VAN BEUSCHOTEN, Professor of Mathematics, Wells College, Aurora, N. Y.

Let the three straight lines be given by the equation

$$y = ax + b, y = cx + d, y = ex + f.$$

The condition that these lines intersect in a common point is given by the vanishing of the determinant,

$$\begin{vmatrix} a & b & 1 \\ c & d & 1 \\ e & f & 1 \end{vmatrix}$$

But the vanishing of this determinant is also the condition that the points (a, b) , (c, d) , (e, f) are colinear.

Also solved by G. B. M. ZERR.

177. Proposed by GEORGE LILLEY, Ph. D., LL. D., University of Oregon, Eugene, Ore.

If two medians of a triangle intersect each other at right angles, the third median will be the hypotenuse of a right triangle, of which the other two will be the sides.

Solution by H. B. PENHOLLOW, DeWitt Clinton High School, New York, N. Y.


Given $\triangle ABC$, medians meeting at O , having $\angle AOB$ a right angle.

From E draw EM perpendicular to AE , meeting AB produced in M . Then $\triangle AEM$ is a right triangle in which AE is one median, $EM = DB$ another median. Also since triangles AEM and AOB are similar, $AE/AO = AM/AB$. But $AE = \frac{3}{2}AO$.

$$\therefore AM = \frac{3}{2}AB.$$

Also OF is median of right triangle AOB .

$$\therefore OF = \frac{1}{2}AB, \text{ or } CF = \frac{3}{2}AB = AM. \quad \text{Q. E. D.}$$



Also solved by P. S. BERG, HENRY HEATON, P. H. PHILBRICK, C. A. LINDEMANN, G. I. HOPKINS, S. E. HARWOOD, J. F. LAWRENCE, T. T. DAVIS, G. B. M. ZERR, and ANNA BENCHOTEN. Professor Penhollow and Miss Benchoten each furnished three solutions.

178. Proposed by JOHN M. ARNOLD, Crompton, R. I.

A cylinder thirty feet long and two feet in diameter is to be placed in a machinery car, the inside dimensions of which are eight feet wide and eight feet high. Find length of the shortest car that will contain it.

No correct solution of this problem has been received.

179. Proposed by ALFRED HUME, C. E., D. Sc., Professor of Mathematics, University of Mississippi, University of Mississippi.

Of all isosceles triangles inscribed in a circle, the equilateral is the maximum and has the maximum perimeter. Prove geometrically.

Solution by the PROPOSER.

Case I. (See Fig. 1.) Vertical angle of isosceles triangle less than 60° .

Let ABC be an inscribed equilateral triangle and ADE any inscribed isosceles triangle with its base DE parallel to BC .